



In partnership with



CUTWORM (TURNIP MOTH) – 28 October 2021

This is the last update for 2021

Page	Contents
2	Introduction
4	Day-degree sum to start of flight activity 2021
5	Pheromone trap captures of turnip moth at Wellesbourne
8	Cutworm forecast
13	Historical information from 2020



Introduction

The 'Cutworm Risk Model'

The cutworm model is a computer program that uses weather data to predict the rate of development of turnip moth eggs and caterpillars. It also predicts the level of rain-induced mortality among the early-instar caterpillars. The cutworm model published by Bowden *et al* (1983) has been programmed into the MORPH decision-support software. This version will be used to produce cutworm forecasts in 2021, with the weather data used to produce the cabbage root fly and carrot fly forecasts.

Reference: Bowden, J., Cochrane, J., Emmett, B. J., Minall, T. E. & Sherlock, P. L. (1983). A survey of cutworm attacks in England and Wales, and a descriptive population model for *Agrotis segetum* (Lepidoptera: Noctuidae). *Annals of Applied Biology* 102 29-47.

Background

“Cutworm” is the name given to caterpillars of certain Noctuid moths, in particular those of the turnip moth *Agrotis segetum*. The name derives from the habit of the older caterpillars of feeding underground, damaging plant roots and stems (including the storage organs that we use for food), sometimes so badly that the plant topples.

The adult moths lay eggs on plants or on pieces of litter and debris in the soil, usually from the end of May or early June. These hatch in around 8-24 days, depending on temperature. The young caterpillars seek out and feed on the aerial parts of plants. In a further 10-20 days, again depending on temperature, the caterpillars go through their second moult, becoming “third instar” caterpillars. It is at this point that they adopt the cutworm habit, becoming subterranean and feeding on roots etc.

Unhatched turnip moth eggs and the older, subterranean cutworms are largely invulnerable to the effects of the weather and insecticides. The two early caterpillar instars differ, however. If there is substantial rainfall (defined as 10 mm or more of rain falling in showers of moderate intensity over a 24-



syngenta.

In partnership with



THE UNIVERSITY OF
WARWICK



hour period) whilst these caterpillars are feeding above ground then this causes high mortality among them. They are also vulnerable to insecticides and irrigation whilst feeding on the foliage.

Crop Susceptibility

Crops differ in their susceptibility to cutworm damage. The most vulnerable are lettuce, leek and red beet. Young lettuce and leek plants are easily bitten through by cutworms, and though beet plants may survive an attack the bulbous root is rendered unmarketable by cutworm feeding. Moderately-susceptible crops include brassicas, carrot, celery, parsnip and sugar beet. The least susceptible of those vulnerable to damage are onion, potato, swede and turnip.




In partnership with



Day-degree sum to start of flight activity 2021

Several years ago, data on trap captures in 2005-7 were used to estimate a day-degree (D°) sum for the start of flight activity at Wellesbourne. This was 340 D° above a base of 7°C from 1 January. The graph below shows accumulated day-degrees for 2021 at Wellesbourne compared with previous years.





In partnership with



Pheromone trap captures of turnip moth at Wellesbourne

Date	Number of moths (2 traps)
21 May	0
25 May	1
28 May	0
1 June	4
4 June	1
8 June	0
11 June	1
15 June	1
18 June	0
22 June	3
25 June	1
29 June	2
2 July	3
6 July	2
9 July	2
13 July	0



In partnership with



Pheromone trap captures of turnip moth at Wellesbourne

Date	Number of moths (2 traps)
16 July	0
20 July	0
23 July	0
27 July	1
30 July	0
3 August	0
6 August	0
10 August	2
13 August	1
17 August	0
20 August	3
24 August	2
27 August	4
31 August	2



In partnership with



Pheromone trap captures of turnip moth at Wellesbourne

Date	Number of moths (2 traps)
3 September	1
7 September	4
10 September	3
14 September	4
17 September	1
21 September	2
24 September	0
28 September	0
1 October	0
5 October	0
8 October	2
12 October	0
Traps removed	



Cutworm forecasts – 2021

The first (male) turnip moth was captured at Wellesbourne between 21st and 25th May. This is slightly earlier than the day-degree forecast. Interestingly although turnip moths live all year round in the UK (and this is what the day-degree forecast assumes), some also arrive as migrant moths, as do diamondback moth and silver Y moth.

The forecasting model will be run (using weather data provided by Plantsystems and funded by Syngenta) from 25th May – based on the capture of the first turnip moth at Wellesbourne. The forecasts indicate that eggs will be hatching from early June. Once eggs start to hatch then rainfall becomes important for the forecast and this is when the forecast outputs will be shown here.

The model has been run for batches of eggs hatching from 1st June. Once eggs start to hatch then rainfall becomes important for the forecast. The table below shows the predicted **percentage survival by 11th August** (the date up to which there are current rainfall data) of caterpillars from eggs hatching each day from 1st June based on rainfall at each site. If there has been no rain at all then 100% of caterpillars are predicted to survive. **Obviously, the amount of rainfall does vary even within a relatively short distance so these are only indications.** The caterpillars are likely to become invulnerable to rainfall and insecticides after about 20 days (when they enter the third instar and burrow in the soil).

This is the last forecast for 2021 as the threat from the first generation of turnip moth is over. There is usually a second generation of turnip moth, but this generation does not appear to be a threat to crops.



In partnership with



	Cornwall	Kent	Suffolk	Warwick	Norfolk	Lincolnshire	Nottinghamshire	Lancashire	Yorkshire	Scotland	Estimated date when no longer affected by rainfall or insecticides
Hatching date	Percentage survival for each hatching date on left hand side										
01-Jun	6	0	0	0	0	0	0	72	21	0	21-Jun
02-Jun	6	0	0	0	0	0	0	72	22	0	22-Jun
03-Jun	6	0	0	0	0	0	0	43	22	0	23-Jun
04-Jun	0	0	0	0	0	0	0	0	0	0	24-Jun
05-Jun	0	0	0	0	0	0	0	4	10	0	25-Jun
06-Jun	0	0	0	0	0	0	0	4	10	0	26-Jun
07-Jun	0	0	0	0	0	0	0	4	10	0	27-Jun
08-Jun	0	0	0	0	0	0	0	4	19	0	28-Jun
09-Jun	0	0	0	0	0	0	0	2	19	0	29-Jun
10-Jun	0	0	0	0	0	0	0	2	19	0	30-Jun
11-Jun	0	0	0	0	0	0	0	2	19	0	01-Jul
12-Jun	0	0	0	0	0	0	0	2	19	0	02-Jul
13-Jun	0	0	0	0	0	0	0	0	14	0	03-Jul
14-Jun	0	0	0	0	0	0	0	0	6	0	04-Jul
15-Jun	0	0	0	0	0	0	0	0	4	0	05-Jul
16-Jun	0	0	0	0	0	0	0	0	0	0	06-Jul
17-Jun	0	0	0	0	0	0	0	0	0	0	07-Jul
18-Jun	0	0	0	0	0	0	0	0	0	0	08-Jul
19-Jun	0	0	0	4	0	0	0	0	0	0	09-Jul
20-Jun	0	0	0	5	0	0	0	0	0	0	10-Jul



In partnership with



	Cornwall	Kent	Suffolk	Warwick	Norfolk	Lincolnshire	Nottinghamshire	Lancashire	Yorkshire	Scotland	Estimated date when no longer affected by rainfall or insecticides
Hatching date	Percentage survival for each hatching date on left hand side										
21-Jun	0	0	0	10	0	0	0	0	0	0	11-Jul
22-Jun	0	0	0	10	0	0	0	0	0	0	12-Jul
23-Jun	0	0	0	10	0	0	0	0	0	0	13-Jul
24-Jun	0	0	0	10	0	0	0	0	0	0	14-Jul
25-Jun	0	0	0	10	0	0	0	0	0	0	15-Jul
26-Jun	0	0	0	12	0	0	0	0	0	0	16-Jul
27-Jun	0	0	0	12	0	0	0	0	0	0	17-Jul
28-Jun	0	1	0	12	0	0	0	0	0	0	18-Jul
29-Jun	0	1	0	25	0	0	0	0	0	0	19-Jul
30-Jun	0	0	0	0	0	0	0	0	0	0	20-Jul
01-Jul	0	2	0	25	0	0	0	0	0	0	21-Jul
02-Jul	0	2	0	25	0	0	0	0	0	0	22-Jul
03-Jul	0	2	0	25	0	0	0	0	0	0	23-Jul
04-Jul	0	2	0	45	0	0	0	0	0	0	24-Jul
05-Jul	0	2	0	69	0	0	0	0	0	0	25-Jul
06-Jul	0	1	0	78	0	0	0	0	0	17	26-Jul
07-Jul	0	2	0	77	36	7	7	5	0	18	27-Jul
08-Jul	0	2	0	62	32	74	2	2	0	0	28-Jul
09-Jul	0	2	0	65	32	74	2	1	0	0	29-Jul
10-Jul	0	2	0	0	0	24	0	1	0	0	30-Jul



In partnership with



	Cornwall	Kent	Suffolk	Warwick	Norfolk	Lincolnshire	Nottinghamshire	Lancashire	Yorkshire	Scotland	Estimated date when no longer affected by rainfall or insecticides
Hatching date	Percentage survival for each hatching date on left hand side										
11-Jul	0	2	0	0	0	23	0	3	0	0	31-Jul
12-Jul	7	0	0	0	0	23	1	3	0	0	01-Aug
13-Jul	7	0	0	0	0	28	2	7	19	0	02-Aug
14-Jul	7	0	0	0	0	19	2	9	19	0	03-Aug
15-Jul	7	0	0	0	0	17	2	8	18	0	04-Aug
16-Jul	4	0	0	0	0	17	0	7	10	0	05-Aug
17-Jul	4	0	0	0	0	17	0	2	2	0	06-Aug
18-Jul	0	0	0	0	0	0	0	0	0	0	07-Aug
19-Jul	0	0	0	0	0	17	0	0	0	0	08-Aug
20-Jul	0	0	0	0	0	17	0	0	0	0	09-Aug
21-Jul	0	0	0	0	0	17	0	0	0	0	10-Aug
22-Jul	0	0	0	0	0	17	0	0	0	0	11-Aug
23-Jul	0	0	0	0	0	17	0	0	0	0	12-Aug
24-Jul	0	0	0	0	0	17	0	0	0	0	13-Aug
25-Jul	0	0	0	0	0	17	0	0	0	0	14-Aug
26-Jul	0	0	0	0	0	17	0	0	0	0	15-Aug
27-Jul	0	0	0	0	0	17	0	0	0	0	16-Aug
28-Jul	0	0	0	0	0	17	0	0	0	0	17-Aug
29-Jul	0	0	0	0	0	18	2	0	0	0	18-Aug
30-Jul	0	0	0	0	0	18	2	0	0	0	19-Aug
31-Jul	0	0	0	47	0	58	8	0	0	0	20-Aug



In partnership with



	Cornwall	Kent	Suffolk	Warwick	Norfolk	Lincolnshire	Nottinghamshire	Lancashire	Yorkshire	Scotland	Estimated date when no longer affected by rainfall or insecticides
Hatching date	Percentage survival for each hatching date on left hand side										
01-Aug	0	0	0	52	0	59	8	0	0	0	21-Aug
02-Aug	0	0	0	56	0	60	8	0	0	0	22-Aug
03-Aug	0	0	0	56	0	60	8	0	0	0	23-Aug
04-Aug	0	0	0	56	0	88	9	0	0	0	24-Aug
05-Aug	0	0	0	56	0	96	9	0	0	0	25-Aug
06-Aug	0	0	0	88	0	98	34	0	1	0	26-Aug
07-Aug	0	0	0	100	0	100	58	0	3	0	27-Aug
08-Aug	37	0	0	100	0	100	68	0	10	0	28-Aug
09-Aug	68	0	0	100	78	100	100	92	71	0	29-Aug
10-Aug	100	100	100	100	99	100	100	94	98	0	30-Aug
11-Aug	100	100	100	100	100	100	100	98	100	98	31-Aug



In partnership with



Historical information from 2020

Date	Numbers of moths captured in 2 pheromone traps at Wellesbourne in 2020 (set up 19 th May)
22 May	1
26 May	7
29 May	6
2 June	7
5 June	5
9 June	6
12 June	2
16 June	2
19 June	1
23 June	0
26 June	1
30 June	0
3 July	1
7 July	0
10 July	0
14 July	0



In partnership with



Date	Numbers of moths captured in 2 pheromone traps at Wellesbourne in 2020 (set up 19 th May)
17 July	0
21 July	3
24 July	2
28 July	1
31 July	0
4 August	0
7 August	1
11 August	0
14 August	1
16 August	5
21 August	25
25 August	25
28 August	8
1 September	13
4 September	4
8 September	0



In partnership with



Date	Numbers of moths captured in 2 pheromone traps at Wellesbourne in 2020 (set up 19 th May)
11 September	0
15 September	0
18 September	0
22 September	1
25 September	0
29 September	4
2 October	0
6 October	0
9 October	0
13 October	0
16 October	0
20 October	2
23 October	1
26 October	1
30 October	2
3 November	1



syngenta.

In partnership with



THE UNIVERSITY OF
WARWICK



Cutworm forecasts - 2020

The model has been run from 20th May – based on the capture of the first turnip moth at Wellesbourne (using weather data provided by Plantsystems and funded by Syngenta). The forecasts indicate that eggs will be hatching from about 2nd June. Once eggs start to hatch then rainfall becomes important for the forecast. The table below shows the predicted **percentage survival by 5th August** (the date up to which there are current rainfall data) of caterpillars from eggs hatching each day from 1st June until 30 July based on rainfall at each site. If there has been no rain at all then 100% of caterpillars are predicted to survive. The caterpillars are likely to become invulnerable to rainfall and insecticides after about 20 days (when they enter the third instar and burrow in the soil). **This is the last forecast for 2020 as the threat from the first generation of turnip moth is over. There is usually a second generation of turnip moth but this generation does not appear to be a threat to crops.**



In partnership with



	Cornwall	Kent	Suffolk	Wellesbourne	Norfolk	South Lincs	Nottingham	Lancashire	York	Scotland	Estimated date when no longer affected by rainfall or insecticides
Percentage survival for each hatching date on left hand side											
01-Jun	0	0	0	0	0	0	0	0	0	0	21-Jun
02-Jun	0	0	0	0	0	0	0	0	0	0	22-Jun
03-Jun	0	0	0	0	0	0	0	0	0	0	23-Jun
04-Jun	0	0	0	0	0	0	0	0	0	0	24-Jun
05-Jun	0	0	0	0	0	0	0	0	0	0	25-Jun
06-Jun	0	0	0	0	0	0	0	0	0	0	26-Jun
07-Jun	0	0	0	0	0	0	0	0	0	0	27-Jun
08-Jun	0	0	0	0	0	0	0	0	0	0	28-Jun
09-Jun	0	0	0	0	0	0	0	0	0	0	29-Jun
10-Jun	0	0	0	0	0	0	0	0	0	0	30-Jun
11-Jun	0	0	0	0	0	0	0	0	0	0	01-Jul
12-Jun	0	0	0	0	0	0	0	0	0	0	02-Jul
13-Jun	0	0	0	0	0	0	0	0	0	0	03-Jul
14-Jun	0	0	0	0	0	0	0	0	0	0	04-Jul
15-Jun	0	0	0	0	0	0	0	0	0	0	05-Jul
16-Jun	0	0	0	0	0	0	0	0	0	0	06-Jul
17-Jun	0	0	0	0	0	0	0	0	0	0	07-Jul
18-Jun	0	0	0	0	0	0	0	0	0	0	08-Jul
19-Jun	0	0	0	0	0	0	0	0	0	0	09-Jul



In partnership with



	Cornwall	Kent	Suffolk	Wellesbourne	Norfolk	South Lincs	Nottingham	Lancashire	York	Scotland	Estimated date when no longer affected by rainfall or insecticides
	Percentage survival for each hatching date on left hand side										
20-Jun	0	0	0	0	0	0	0	0	0	0	10-Jul
21-Jun	0	0	0	0	0	0	0	0	0	0	11-Jul
22-Jun	0	0	0	0	0	0	0	0	0	0	12-Jul
23-Jun	0	0	0	0	0	0	0	0	0	0	13-Jul
24-Jun	0	0	0	0	0	0	0	0	0	0	14-Jul
25-Jun	0	0	0	0	0	0	0	0	0	0	15-Jul
26-Jun	0	0	0	0	0	0	0	0	0	0	16-Jul
27-Jun	0	0	0	0	0	0	0	0	0	0	17-Jul
28-Jun	0	0	0	0	0	0	0	0	0	0	18-Jul
29-Jun	0	0	0	0	0	0	0	0	0	0	19-Jul
30-Jun	0	0	0	0	0	0	0	0	0	0	20-Jul
1-Jul	0	0	0	0	0	0	0	0	0	0	21-Jul
2-Jul	0	0	0	0	0	0	0	0	0	0	22-Jul
3-Jul	0	0	0	0	0	0	0	0	0	0	23-Jul
4-Jul	0	0	0	0	0	0	0	0	0	0	24-Jul
5-Jul	0	0	0	0	0	0	0	0	0	0	25-Jul
6-Jul	0	0	0	0	0	0	0	0	0	0	26-Jul
7-Jul	0	0	0	0	0	0	0	0	0	0	27-Jul
8-Jul	0	0	0	0	0	0	0	0	0	0	28-Jul



In partnership with



	Cornwall	Kent	Suffolk	Wellesbourne	Norfolk	South Lincs	Nottingham	Lancashire	York	Scotland	Estimated date when no longer affected by rainfall or insecticides
	Percentage survival for each hatching date on left hand side										
9-Jul	0	0	0	0	0	0	0	0	0	0	29-Jul
10-Jul	0	0	0	0	0	0	0	0	0	0	30-Jul
11-Jul	0	0	0	0	0	0	0	0	0	0	31-Jul
12-Jul	0	0	0	0	0	0	0	0	24	0	01-Aug
13-Jul	0	0	0	0	0	0	0	0	24	0	02-Aug
14-Jul	0	0	0	0	0	0	0	0	24	0	03-Aug
15-Jul	0	0	0	6	0	0	0	0	24	0	04-Aug
16-Jul	0	0	55	6	0	0	0	0	24	0	05-Aug
17-Jul	0	0	55	6	0	0	0	0	24	0	06-Aug
18-Jul	0	0	55	6	0	0	0	0	24	0	07-Aug
19-Jul	0	0	55	6	0	0	0	0	24	0	08-Aug
20-Jul	0	0	55	32	0	0	0	0	26	0	09-Aug
21-Jul	0	16	59	64	0	0	0	0	26	0	10-Aug
22-Jul	0	16	59	64	0	0	0	0	26	0	11-Aug
23-Jul	0	16	59	64	0	0	0	0	26	0	12-Aug
24-Jul	0	16	59	64	0	0	0	0	26	0	13-Aug
25-Jul	0	16	59	64	0	0	0	0	32	0	14-Aug
26-Jul	0	16	59	64	0	0	0	0	34	0	15-Aug
27-Jul	26	88	87	74	30	0	50	0	40	0	16-Aug
28-Jul	36	90	89	74	30	2	50	0	44	0	17-Aug
29-Jul	100	100	100	100	100	100	100	96	100	43	18-Aug
30-Jul	100	100	100	100	100	100	100	96	100	43	19-Aug



In partnership with



Day-degree sum to start of flight activity 2020

Several years ago, data on trap captures in 2005-7 were used to estimate a day-degree (D°) sum for the start of flight activity at Wellesbourne. This was 340 D° above a base of 7°C from 1 January. The graph below shows accumulated day-degrees for 2020 at Wellesbourne compared with previous years. **Flight was predicted to start on 20th May.**

