



2210/2025

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

syngenta.

Syngenta UK Limited,

Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY

Tel: +44 (0) 1223 883400

**In case of toxic or transport emergency ring +44 (0) 1484 538444 anytime**

**GROUP 7 FUNGICIDE**

Product registration number:

MAPP 19174

UFI: D94A-90KU-500M-S16T



APROVIA® Plus is an emulsifiable concentrate containing 100 g/l (10.2% w/w) benzovindiflupyr.

Provides control of *Mycosphaerella graminicola*, Brown rust and Yellow rust and moderate control of *Septoria nodorum* in winter and spring wheat, control of Net blotch, *Ramularia collo-cygni* and Brown rust and moderate control of *Rhynchosporium secalis* on winter and spring barley, control of *Septoria spp.*, Brown Rust and Yellow Rust and moderate control of *Rhynchosporium secalis* on triticale, control of Brown rust and *Rhynchosporium secalis* on rye, control of *Puccinia spp.* on onion, garlic, shallot, leeks and salad onions (spring onion, green onion and welsh onion).

PROTECT FROM FROST

SHAKE WELL BEFORE USE



The  
Voluntary  
Initiative

This product label is compliant with the CPA Voluntary Initiative (VI) guidance.

L1151327 GBRI/04E PPE 4252234

**3 litres**

Product names marked © or ™, the ALLIANCE FRAME the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company



**APROVIA® Plus**

Emulsifiable concentrate containing 100 g/l (10.2% w/w) benzovindiflupyr.

**Danger**

**Harmful if swallowed or inhaled.**

**Causes serious eye damage.**

**May cause an allergic skin reaction.**

**May cause respiratory irritation.**

**Very toxic to aquatic life with long lasting effects.**

Avoid breathing mist or vapours.

Wash skin thoroughly after handling.

Wear protective gloves/eye protection/face protection.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

Call a POISON CENTRE/doctor if you feel unwell.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician. Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment comply with the instructions for use.**

**MAPP NO. 19174 UFI: D94A-90KU-500M-S16T**

**IMPORTANT INFORMATION**

**FOR USE ONLY AS A PROFESSIONAL FUNGICIDE**

<b>Crops</b>	Winter and spring wheat, rye and triticale.	Winter and spring barley.	Onion, garlic, shallot, leeks and salad onions
<b>Max. individual dose (litres/hectare/crop)</b>	0.75		0.50
<b>Max. no. of treatments (per crop)</b>	1		
<b>Latest time of application</b>	Up to and including anthesis complete (GS 69).	Up to and including complete ear emergence (GS 59).	Up to and including leaves bent over 50% of plants (GS 48)
<b>Aquatic buffer zone distance (metres)</b>	6		

**Other specific restrictions:**

1. This product must not be applied via hand-held equipment.
2. No more than two applications of products containing SDH inhibitors must be applied to any cereal crop.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## **SAFETY PRECAUTIONS**

### **(a) Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**OPERATORS MUST WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS), SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD)** when handling the concentrate.

However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

**WASH SPLASHES** from skin immediately.

**WASH HANDS AND EXPOSED SKIN** before meals and after work.

### **(b) Environmental protection**

To protect aquatic organisms, respect an unsprayed buffer zone to surface water bodies as specified for the crop.

**HORIZONTAL BOOM SPRAYERS MUST BE FITTED WITH THREE STAR DRIFT REDUCTION TECHNOLOGY** for all uses. Low drift spraying equipment must be operated according to the specific conditions stated in the official three star rating for that equipment as published on HSE Chemicals Regulation Division's website. Maintain three star operating conditions until 30 m

from the top of the bank of any surface water bodies.

**DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within the distance specified for the crop to the top of the bank of a static or flowing water body, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water. **NOTE: BUFFER ZONES OF MORE THAN 5 M CANNOT BE REDUCED UNDER THE LOCAL ENVIRONMENT RISK ASSESSMENT FOR PESTICIDES (LERAP) SCHEME.**



The statutory buffer zone must be maintained and the distance recorded in Section A of the LERAP record form. The LERAP record form must be kept available for three years.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### **(c) Storage and disposal**

**RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. **KEEP IN ORIGINAL CONTAINER**, tightly closed, in a safe place. **EMPTY CONTAINER COMPLETELY** and dispose of safely.

## **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product. Consult processors before using on crops of garlic, shallot, bulb onion and salad onions grown for processing.

## **GENERAL INFORMATION**

Benzovindiflupyr is an orthosubstituted pyrazole carboxamide fungicide belonging to the sub-class of the benzonorbornenes. Benzovindiflupyr is an SDH inhibitor (FRAC group #7 carboxamides). Benzovindiflupyr is predominantly a protectant substance.

APROVIA® Plus is best used as a protectant treatment or in the earliest stages of disease development.

## **DISEASES CONTROLLED**

### **Winter and Spring Wheat**

Leaf spot (*Mycosphaerella graminicola*)

Glume blotch (*Septoria nodorum*) [Moderate control]

Yellow rust (*Puccinia striiformis*)

Brown rust (*Puccinia recondita*)

### **Winter and Spring Barley**

Net blotch (*Pyrenophora teres*)

Leaf blotch (*Rhynchosporium secalis*) [Moderate control]

Brown rust (*Puccinia hordei*)

*Ramularia collo-cygni*

### **Triticale**

Yellow rust (*Puccinia striiformis*)

Brown rust (*Puccinia recondita*)

*Septoria* spp.

Leaf blotch (*Rhynchosporium secalis*) [Moderate control]

### **Rye**

Brown rust (*Puccinia recondita*)

Leaf blotch (*Rhynchosporium secalis*)

### **Onion, garlic, shallot, leeks and salad onions**

*Puccinia* spp.

## **RESISTANCE MANAGEMENT**

Use APROVIA Plus as part of an Integrated Crop Management (ICM) strategy incorporating other methods of control, including where appropriate other fungicides with a different mode of action.

You must not apply more than two foliar applications of products containing SDH inhibitors to any cereal crop.

Disease control may be reduced if strains of pathogens less sensitive to APROVIA Plus develop.

On cereal crops, APROVIA Plus must always be used in mixture with another product, recommended for control of the same target disease that contains a fungicide from a different cross resistance group and is applied at a dose that will give robust control.

Users should refer to current FRAG-UK guidelines for SDHI compounds.

## **CROP SPECIFIC INFORMATION**

### **Crops and growing conditions**

APROVIA Plus can be used on all varieties of winter and spring wheat, winter and spring barley, rye, triticale, onion, garlic, shallots, leeks and salad onions. Apply APROVIA Plus under good growing conditions with adequate soil moisture. Avoid poor growing conditions which may give less reliable results.

## **Timing**

Always inspect crops to assess disease development immediately before spraying. Best results will be achieved from applications made as a protectant treatment or in the earliest stages of disease development following a disease risk assessment or the use of appropriate decision support systems.

## **Rates of use**

Apply APROVIA Plus at 0.75 litres per hectare.

Onion, garlic, shallot, leeks and salad onions: Apply APROVIA PLUS at 0.50 litres per hectare.

## **FOLLOWING CROPS**

There are no restrictions on succeeding crops in a normal rotation.

## **MIXING AND SPRAYING**

### **Mixing Procedure**

Make sure the sprayer is set to give an even application at the correct volume. Fill the spray tank with half the required volume of water and begin agitation. Add the required amount of APROVIA Plus to the spray tank and allow to disperse before adding any other product. Add the rest of the water

and continue to agitate the mixture thoroughly. Always agitate during spraying.

### **Spray Volume and Application**

Apply APROVIA Plus in a recommended 100 - 400 litres of water per hectare through conventional crop spraying equipment. The higher spray volumes are recommended where the crop is dense or disease pressure/risk is high to ensure good penetration to the lower leaves and stem bases. Disease control may be compromised by reducing water volumes, where good spray coverage is difficult to achieve. Apply APROVIA PLUS in a recommended 300 - 600 litres of water per hectare for bulb vegetables. A spray pressure of 2-3 bars is recommended. Effectiveness using three star drift reduction technology may be reduced.

### **After Spraying**

Thoroughly wash out sprayer according to manufacturer's guidelines and dispose of washing and clean containers according to DEFRA Code of Practice and local water authority guidelines.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier.

**APROVIA® Plus**

Emulsifiable concentrate containing 100 g/l (10.2% w/w) benzovindiflupyr.

**Danger**

**Harmful if swallowed or inhaled.**

**Causes serious eye damage.**

**May cause an allergic skin reaction.**

**May cause respiratory irritation.**

**Very toxic to aquatic life with long lasting effects.**

Avoid breathing mist or vapours.

Wash skin thoroughly after handling.

Wear protective gloves/eye protection/face protection.

IF INHALED: Remove person to fresh air and keep comfortable for breathing.

Call a POISON CENTRE/doctor if you feel unwell.


IF IN EYES: Rinse cautiously with water for several minutes.

Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTRE or doctor/physician. Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

**To avoid risks to human health and the environment comply with the instructions for use.**

**MAPP NO. 19174 UFI: D94A-90KU-500M-S16T**

**IMPORTANT INFORMATION**

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

<b>Crops</b>	Winter and spring wheat, rye and triticale.	Winter and spring barley.	Onion, garlic, shallot, leeks and salad onions
<b>Max. individual dose (litres/hectare/crop)</b>	0.75		0.50
<b>Max. no. of treatments (per crop)</b>	1		
<b>Latest time of application</b>	Up to and including anthesis complete (GS 69).	Up to and including complete ear emergence (GS 59).	Up to and including leaves bent over 50% of plants (GS 48)
<b>Aquatic buffer zone distance (metres)</b>	6		

**Other specific restrictions:**

1. This product must not be applied via hand-held equipment.
2. No more than two applications of products containing SDH inhibitors must be applied to any cereal crop.

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**