



syngenta

BERET GOLD contains fludioxonil and controls seed-borne diseases on winter and spring crops of wheat, barley, oats, rye, triticale and also soil-borne bunt.

SAFETY PRECAUTIONS

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling treated seed or contaminated surfaces.

THIS SEED HAS BEEN CHEMICALLY TREATED.

DO NOT HANDLE seed unnecessarily.

DO NOT USE TREATED SEED as food or feed.

DO NOT RE-USE SACKS OR CONTAINERS THAT HAVE CONTAINED TREATED SEED for food or feed.

KEEP TREATED SEED SECURE from people, domestic stock/pets and wildlife at all times during storage and use.

WASH HANDS AND EXPOSED SKIN before meals and after work.

BURY OR REMOVE SPILLAGES.

HARMFUL TO GAME AND WILDLIFE.

Treated seed should not be left on the soil surface. Bury spillages.
DO NOT APPLY TREATED SEED FROM THE AIR.

CAUTION

BERET GOLD may reduce the flow rate of seed in a seed drill. Recalibrate drill with BERET GOLD treated seed before drilling.

NOTES

1. Safe Handling of Treated Seed.
Avoid skin contact with treated seed and dust during all drilling operations.
Launder coveralls daily.

2. Drilling
Check drill calibration before drilling

for each batch of seed to ensure an accurate drilling rate. Avoid adverse seedbed conditions and deep or shallow drilling which may adversely affect crop establishment and reduce the level of pest control.

3. Storage

Seed should be stored in a cool, dry, well ventilated building and be drilled as soon as possible after treatment. Drill during the season of treatment.

4. Seed Spillages

In case of seed spillage, clean up as much as possible into the related seed sack and re-use the clean seed. Bury the remainder completely.

Syngenta UK Limited, Jealott's Hill International Research Centre, Bracknell, Berkshire, Tel +44 (0) 1223 883400.
In case of toxic or transport emergency ring +44 (0) 1484 538444 any time

BERET GOLD® is a Registered Trademark of a Syngenta Group Company. © Syngenta AG 2026.

