

Powered by
Syngenta Vegetable Seeds

DISCOVER OUR VEGETABLE VARIETIES



syngenta®



Contents

| | |
|---------------------|---|
| Introduction | 4 |
| Meet the Team | 5 |
| Clubroot Solutions | 6 |
| FarMore® Technology | 9 |

Discover our Varieties

| | |
|------------------------|----|
| Broccoli | 10 |
| Brussels Sprouts | 14 |
| Borecole | 20 |
| Cabbage | 21 |
| Cauliflower | 29 |
| Wholehead Lettuce | 39 |
| Babyleaf | 48 |
| Onions | 57 |
| Squash | 60 |
| Table of Abbreviations | 63 |
| Levels of Resistance | 64 |
| Notes | 66 |

Breeding to Meet the Next Green Revolution

Vegetable crop genetics offer the first step to tackle many of the difficult challenges currently impacting on the profitability of UK growers. Syngenta is at the forefront of delivering varieties developed to meet growers' changing needs.

Your variety selection will have a major influence on business performance at all levels, from practical agronomy decisions, to resolving labour shortages, managing increasing costs, coping with changing climate, creating innovative markets and, ultimately, satisfying the end-consumer.

Now, with the industry's extra focus on delivering a reduction in carbon emissions, variety choice will be crucial in driving towards net zero targets.

For our specialist vegetable and salad crop variety breeders, high yield remains a key objective for any new development. However, greater focus on the traits for efficiency of production and marketability could lead to improved overall profitability and business sustainability.

Working with our dedicated UK field team to tailor variety characteristics best suited to a farm's individual situation can deliver better returns for the current season and justify investment in your business for the future.

Making more efficient use of every input will be crucial for lowering costs and reducing environmental impacts of production. With escalating fertiliser costs, for example, varieties available now that perform well – or in some cases better – under a low-nutrition regime offer real benefits. Drought and heat tolerance will be essential variety attributes to make more effective use of water resources in the future.

Breeding varieties with the resilience to cope with climate extremes better assures consistent and predictable production for you and your customers. Syngenta's pan-European and global reach gives access to vegetable crop

genetics and experience that will be invaluable for UK growers to adapt.

Today's Syngenta variety selection could enable UK growers to supply an extended season of home-grown produce in demand from customers – reducing costs, challenges and carbon footprint of long-distance transport.

With labour availability now a major issue for growers across Europe, which has been amplified in the UK since Brexit, means mechanical harvesting and processing requires significant investment from growers. Varieties bred with the robustness and uniformity to suit mechanical harvesting systems are essential for the exciting technology to be effectively utilised and to justify the investment.

Furthermore, Syngenta's development of vegetable and salad varieties that require less trimming and processing helps to save time and cost on packing lines, as well as reducing waste right through to the retail shelf and enhanced end-consumer satisfaction.

For the future, varieties that will perform consistently well in Integrated Pest Management (IPM) systems will become increasingly important. Interpreting responses to biological, biopesticide and biostimulant inputs is only possible through extensive trials and specialist understanding of plants' genetic potential.

The continued ability of UK vegetable and salad growers to adapt to new challenges is ever more important. Syngenta seed breeders and our UK field teams are committed to developing and supporting the exciting and innovative varieties highlighted in this catalogue, that will meet your needs now and in the future.

Meet the Team



Kris Goen
Seeds Commercial Manager
UKI & Nordics
T: +32 498.85.80.63
E: kris.goen@syngenta.com



Nigel Kingston
Technical Sales Representative
(Onion, Broccoli and Cabbage)
T: 07711 655526
E: nigel.kingston@syngenta.com



Harry Twinberrow
Technical Sales Representative
(Brussels Sprouts and Kale)
T: 07385 466591
E: harry.twinberrow@syngenta.com



Louis Stokes
Technical Sales Representative
(Cauliflower)
T: 07776 994697
E: louis.stokes@syngenta.com



Rosie Frost
Technical Sales Representative
(Leafy crops, Courgette and Pumpkin)
T: 07776 997416
E: rosemary.frost@syngenta.com



Luke Duffy
Junior Breeding Trials Specialist
(Leafy)
T: 07484 926799
E: luke.duffy@syngenta.com



Sue Hutchinson
Marketing Operations Lead
T: 07484 908946
E: susan.hutchinson@syngenta.com



Henrik Pedersen
Technical Sales Representative –
Nordics
T: +45 20151763
E: henrik.pedersen@syngenta.com



Sanna Hasila
Technical Sales Representative –
Nordics
T: +35 8400 285 783
E: sanna.hasila@syngenta.com



Rebecca White
Vegetable Business Manager
(Crop Protection)
T: 07894 392984
E: rebecca.white@syngenta.com



Customer Services
T: 0845 2660014
T: +31 (0) 228 366 306
E: VegetablesCSDesk.UK@syngenta.com



Clubroot
(Plasmodiophora brassicae) is a serious disease affecting Brassica. An estimated 10% of the total cultured area worldwide contracts the disease.



Syngenta has invested in years of breeding to offer a high level of resistance in their Brassica varieties.

Syngenta has pioneered robust and reliable clubroot resistance in a range of key brassica varieties since 2005



Clubroot symptoms

Clubroot is a soil-borne fungal disease which attacks the roots of Brassica crops. It is considered as one of the most economically important diseases of cultivated crucifers. Roots affected by clubroot are swollen and distorted. The damage caused to the roots causes crops to be stunted and, in most cases, there is a reduction in yield. The pathogen survives in the soil for up to 15 years in the form of resting spores released from decayed galls.



Syngenta's solutions

The potential of cultural practices to reduce crop losses due to clubroot are limited and chemical treatments to control the fungus are either banned, due to environmental regulations, or are not cost-effective. The best way to combat clubroot is through the breeding of resistant varieties. Syngenta has succeeded, after many years of breeding, to introduce a high level of resistance in varieties of cabbage (Chinese, White & Green), Brussels sprouts, broccoli and cauliflower. Syngenta will, over the coming years, be introducing the resistance across the Brassica range.

We currently have the following varieties with clubroot resistance:

| BROCCOLI | BRUSSELS SPROUTS | SAVOY CABBAGE | WHITE CABBAGE | CAULIFLOWER |
|--------------------------------|------------------|----------------------------|---------------|-------------|
| MONCLANO | CRISPUS | CORDESA | KILASTOR | CLAPTON |
| NPI <small>COMING SOON</small> | CRYPTUS | CORDOBA | KILAZOL | CLARIFY |
| | | <small>NEW</small> CORRIPA | KILACEES | CLARINA |
| | | <small>NEW</small> CORADI | | CLEOZIL |

LOSSES FROM CLUBROOT

Growing cost per ha

£3,960*

+ loss of income per ha

£5,720

*Excludes rent.
Source: John Nix 2022



Syngenta continues to develop a new generation of clubroot resistant varieties providing protection against an even wider range of clubroot strains.

Clubroot resistant varieties reduce the risk of plant loss and enable growers to manage cropping areas with known levels of infection.



What is FARMORE® Technology?

FarMore® Technology is the first comprehensive combination of **separately-registered seed protection products, proprietary application technologies and dedicated seed treatment services** that maximise vegetable production value by enhancing performance and quality.



What is APRON® XL and what does it do?

APRON XL is a modern systemic seed treatment fungicide. APRON XL is specially developed for seed treatment and contains 35% mefenoxam, the biologically most active isomer of the compound metalaxyl.

APRON XL controls seed and soil-borne fungi and downy mildew. The product is quickly absorbed by the seed and spreads through the plant after germination. This protects the seed and the seedling at an early stage, resulting in a fast development and a uniform crop. APRON XL is effective in damp, colder conditions that are ideal for *Pythium*, and warmer, humid conditions ideal for *Phytophthora* and downy mildew. APRON XL is registered in a wide range of vegetable crops.



What is MAXIM® 480FS and what does it do?

MAXIM 480FS is a special seed treatment that is effective against a broad spectrum of seed and soil-borne diseases in a wide range of vegetable crops. Active ingredient fludioxonil is a contact fungicide that penetrates the seed surface and coats the seed, providing long-lasting protection around the young seedling and combating diseases such as *Alternaria*, *Phoma* and *Fusarium*. It has excellent activity at low rates, has a positive effect on germination capacity and plant vigour, and is suitable as a mixing partner with other seed treatments.



DISCOVER OUR CAULIFLOWER VARIETIES

“Working with growers and breeders, to ensure we deliver new cauliflower varieties, that can reliably produce product in all conditions throughout the year. This really adds value for our customers and can only be achieved with the growers and breeders at the heart of our product development.”



MEET THE EXPERT

Louis Stokes
Technical Crop Advisor
(Cauliflower)

T: 07776 994697

E: louis.stokes@syngenta.com

Our Cauliflowers with High Resistance to Clubroot




Clapton

- Clubroot resistant summer and autumn cauliflower
- Used for fresh market and processing
- Heavy curds/dense florets

Normal programme

Key: ● Plant ● Harvest




Clarify

- Clubroot resistant summer production
- High vigour plants
- Suitable for fresh market and processing

Normal programme

Key: ● Plant ● Harvest




Clarina

- Summer production
- High vigour
- For fresh market and processing

Normal programme

Key: ● Plant ● Harvest




SGC2101
 Cleozil

- 75-80 growing days
- Very healthy plant
- Uniform curds
- Good frame size

Nitrogen advice: More nitrogen at transplanting

Normal programme

Key: ● Plant ● Harvest



Our Standard Cauliflower Varieties




 **Almagro**

- 85-90 growing days
- Firm curd, reliable in quality
- Very good self-protection
- Nice, dense white curds
- Good vigour and plant health
- Strong field standing ability
- Easy to harvest and bags well

Normal programme

Key: ● Plant ● Harvest



 **Andromeda**

- Excellent shelf-life
- High planting density possible
- Vigorous crop
- High uniform harvest
- Can be over-winter sown
- Maintains colour in the sun
- Open habit, making it visible and easier to harvest

Normal programme

Key: ● Plant ● Harvest



Over-winter programme




 **Electra**

- Good self protection curd
- Uniform, quality harvest
- Good density and firmness
- Good, healthy plant
- Nice white colour

Normal programme

Key: ● Plant ● Harvest




 **Koten**

- Excellent, self protection
- Uniform harvest
- Heavy, firm, white curd
- Good vigour

Normal programme

Key: ● Plant ● Harvest




Our Coloured Cauliflower Varieties



SGC8209 Clementine

- Uniform harvest
- Very good vigour
- Dark orange colour
- Easy to grow variety


Normal programme Key: ● Plant ● Harvest



Graffiti

- Large, dense curd
- Good field standing
- Attractive, dark purple colour
- Strong, vigorous and well formed


Normal programme Key: ● Plant ● Harvest



SGC8001 Greenstorm

- 65-75 growing days
- Vibrant green curds
- Good against bracting
- Medium cycle maturing variety
- Summer and autumn
- Unique, short growing cycle

Normal programme Key: ● Plant ● Harvest



Our Long Cycle Winter Cauliflower Varieties




SGC4049
Kashen

- Early winter production
- Heavy white curds
- Good vigour and healthiness

Normal programme

Key: ● Plant ● Harvest




LEOCEN

- Long cycle variety
- Plant habit comparable to Lestren
- Matures 10 days later than Lestren

Normal programme

Key: ● Plant ● Harvest




SGC5073
Lirren

- White curd
- Great variety for Christmas
- Uniform harvest

Normal programme

Key: ● Plant ● Harvest




SGC4044
Akinen

- Excellent self protection
- Good vigour
- Firm, heavy white curd
- Performs well across the UK

Mid-winter programme – Cornwall

Key: ● Plant ● Harvest



Our Very Long Cycle Winter Cauliflower Varieties




 **Amonen**

- Late alpen variety
- Dark green foliage
- Good wrap

Mid-winter programme – Cornwall

Key: ● Plant ● Harvest




 **Caylen**

- Excellent self-protection and frost tolerance
- Good uniformity and dense, solid curd
- Proven reliable performance

Early winter programme

Key: ● Plant ● Harvest




NEW

SGC5086
 **Adrien**

- Healthy, vigorous plant
- Resistant to *Mycosphaerella*
- Good density
- Good and easy to see SEP

Mid-winter programme – Cornwall

Key: ● Plant ● Harvest




NEW

 **Clemen**

- Consistent high percentage cut out
- Vigorous and healthy plant frame
- Presents itself well when mature – easy to harvest

Normal programme

Key: ● Plant ● Harvest



Our Very Long Cycle Winter Cauliflower Varieties



SGC5044
Darifen

- Uniform maturity
- Excellent vigour
- Good quality curds

Mid-winter programme – Cornwall

Key: ● Plant ● Harvest



Mid-winter programme – Lincolnshire



SGC5033
Dragen

- Good base health
- Good curd quality

Mid-winter programme – Cornwall

Key: ● Plant ● Harvest



Mid-winter programme – Lincolnshire




Delon

- Flexible, sequential sowings can be made to spread maturity
- Excellent storage potential for added security
- Intermediate resistance to *Mycosphaerella*


Normal programme

Key: ● Plant ● Harvest



VARIETY FIT OVERVIEW CAULIFLOWER


Key: ● Plant ● Harvest



*Clubroot resistant

VARIETY FIT OVERVIEW CAULIFLOWER

Key: ● Plant ● Harvest





Levels of Resistance

Two levels of resistance are defined:

High/standard resistance (HR*)

Plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. These plant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.

Moderate/intermediate resistance (IR*)

Plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to high/standard resistant varieties. Moderately/intermediately resistant plant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar

environmental conditions and/or pest or pathogen pressure. Susceptibility is the inability of a plant variety to restrict the growth and development of a specified pest or pathogen.

The Vegetable Section of ISF recommends, as it pertains to biotic stress, that its members use the terms immunity, high/standard or moderate/intermediate resistance and susceptibility and to avoid the term tolerance in communications with their customers.

Tolerance is the ability of a plant variety to endure abiotic stress without serious consequences for growth, appearance and yield. Vegetable companies will continue to use tolerance for abiotic stress.

Fungal disease – what's it worth?

1% reduction in pack out

£90/ha*

*Excludes grading and transport.
Approximate figures based on Brussels sprouts output – 20 t/ha @ £450/t.

Growers can rely on good levels of resistance against *Mycosphaerella* (Ringspot) in our varieties and we go further in our ambition to deliver solutions.

We introduced the first Savoy cabbage varieties to the market to combine resistance against *Mycosphaerella*, white blister and clubroot. With unpredictable seasons, this combined resistance against several fungal diseases in one variety helps assure quality whatever the season.

Customer Services

Tel: 0845 2660014

Tel: +31 (0) 228 366 306

Email: VegetablesCSDesk.UK@syngenta.com

Technical enquiries

Tel: 0800 169 6058

Email: customer.services@syngenta.com

Syngenta UK Ltd

CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE

Tel: 01223 883400 Fax: 01223 882195

Website: www.syngenta.co.uk

APRON® XL and MAXIM® 480FS are Registered Trademarks of a Syngenta Group Company. APRON XL (MAPP 14654) contains metalaxyl-M and MAXIM 480FS (MAPP 16725) contains fludioxonil. All other brand names used are Trademarks of other manufacturers in which proprietary rights may exist. Use plant protection products safely. **Always read the label and product information before use. For further information including warning phrases and symbols refer to the website www.syngenta.co.uk**

Syngenta UK Ltd, Registered in England No. 849037, CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE.
Tel: 01223 883400 Fax: 01223 882195 Technical Enquiries Tel: 0800 169 6058
Email: customer.services@syngenta.com ©Syngenta Feb 2022. 11674.

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure.

All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge.

Syngenta Seeds Vegetables uses highly elaborate analytical methods to verify specific variety resistances. Specificity of pests or pathogens may vary over time and depends on environmental factors.

In order to maximise the efficiency of a resistance, it is highly recommended to mix different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management.

The Syngenta resistance against Clubroot is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields.

Genetic resistance is only one of the tools to manage Clubroot. Cultural measures such as liming, use of fertilisers with high percentage of calcium, proper drainage, good crop hygiene management are several important components of an integrated approach to manage the disease.

We always recommend to first execute small variety trials before starting commercial production of a new variety.

Spinach leaf spots can be caused by many different fungus; i.e. *Peronospora effusa*, *Stemphylium* spp., *Cladosporium variabile*, *Colletotrichum dematium* which are not always monitored by EU authorities.

Syngenta identified high resistance (HR) in our genetic to at least one *Stemphylium* specie that we identified & isolated from many leafspot samples over the last few years & in different EU countries.

The latest International Seed Federation (I.S.F.) terms and definitions describing the reaction of plants to pests and pathogens and to abiotic stresses for the vegetable seed industry are hereby incorporated by reference. The meaning of such terms in any related statement made by Syngenta shall be as provided by the I.S.F. If Syngenta adopts a proper term to define the reaction of plants to pests and pathogens and to abiotic stresses, Syngenta shall inform the customers of such term and of its definition.

All data in this brochure are intended for general guidance only and the user should apply it in accordance with their own knowledge and experience of local conditions. In case of doubt we recommend that a small scale production trial be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.