



# Amphore<sup>®</sup> Plus

syngenta<sup>®</sup>

**GROUP 40/3 FUNGICIDES**

**F**



**(b) Environmental protection**

To protect aquatic organisms, respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

**DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a horizontal boom sprayer or broadcast air assisted sprayer either a LERAP must be carried out in accordance with CRD published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads. **RISK TO NON-TARGET INSECTS OR OTHER ARTHROPODS.** See Directions for Use.

**(c) Storage and disposal**

**KEEP IN ORIGINAL CONTAINER**, tightly closed, in a safe place **DO NOT RE-USE CONTAINER** for any purpose. **RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

**Product reg. no:** MAPP 16327 **UFI:** CFM2-NOQ7-HOOG-T6UU

AMPHORE<sup>®</sup> Plus is a suspension concentrate containing 250 g/l mandipropamid and 250 g/l difenoconazole.

For the control of Foliar blight (*Phytophthora infestans*) and moderate control of Early blight (*Alternaria solani*, *Alternaria alternata*) in potatoes.

*The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.*

**In case of toxic or transport emergency ring**  
**+44 (0)1484 538444 any time**

Syngenta UK Limited  
Jealott's Hill International Research Centre,  
Bracknell, Berkshire, RG42 6EY  
Tel: +44 (0) 1223 883400



This product label is compliant with the CPA Voluntary Initiative (VI) guidance.

**SHAKE WELL BEFORE USE.**  
**PROTECT FROM FROST.**

**SAFETY PRECAUTIONS**

**(a) Operator protection**

**WASH HANDS AND EXPOSED SKIN** before eating, drinking and after work.  
**IF YOU FEEL UNWELL**, seek medical advice (show label where possible).



L1127295 GBRI/10C PPE 4219319

**5 litres**

AMPHORE® Plus is a suspension concentrate containing 250 g/l mandipropamid and 250 g/l difenoconazole

### Warning

**May be harmful if swallowed**

**Very toxic to aquatic life with long lasting effects.**

If medical advice is needed, have product container or label at hand.

Keep out of reach of children.

Wash hands thoroughly after handling.

Do not eat, drink or smoke when using this product.

Avoid release to the environment.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

**To avoid risks to human health and the environment, comply with the instructions for use.**



**Product reg. no:** MAPP 16327 **UFI:** CFM2-N0Q7-H00G-T6UU

### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

For use on potatoes

Maximum Individual Dose:	0.6 litres product per hectare
Maximum Number of Treatments:	Three per crop
Latest Time of Application:	3 days before harvest

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS**

This leaflet is part of the approved Product Label.

## **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

Avoid spraying within 5m of the field boundary to reduce effects on non-target insects or other arthropods.

These buffer distances should be measured from the field boundary, which for the purposes of this labelling is defined as from the edge of non-cropped land (i.e. land taken permanently out of agricultural production, including the 1-2 m strips adjacent to hedgerows and watercourses established under the Single Payment Scheme). Cropped land includes managed buffer strips (e.g. grass strips, wild flower margins and conservation headlands), but since these are usually set up as havens for wildlife it is best practice to minimise spray drift onto them

AMPHORE PLUS can be used on all varieties of potatoes including seed potatoes.

## **PROPERTIES OF AMPHORE PLUS**

AMPHORE PLUS is a foliar fungicide for the preventative moderate control of Early blight (*Alternaria solani*, *Alternaria alternata*) and control of Late blight (*Phytophthora infestans*) in potatoes and is highly active against spore germination. Uptake into the leaf tissue assures good translaminar activity and inhibits mycelial growth during the incubation period.

## **COMPANY ADVISORY INFORMATION**

As part of our Product Stewardship policy, Syngenta UK Limited recommends the following precautions should also be observed:

Wear appropriate clothing – cotton overall and rubber gloves, when handling the concentrate.

Since the occurrence of resistance cannot be forecast, neither Syngenta UK Limited nor its distributors can accept responsibility for any loss or damage to crops caused by the failure of AMPHORE PLUS to control resistant strains.

## **DISEASE CONTROLLED**

For the control of Foliar blight (*Phytophthora infestans*) and moderate control of Early blight (*Alternaria solani*, *Alternaria alternata*) in potatoes.

## **BACKGROUND TO POTATO BLIGHT CONTROL**

Late blight (*Phytophthora infestans*) is potentially a devastating disease of potatoes. In commercial production, a season long **disease prevention** policy is essential. First of all ensure that other control methods are being satisfactorily applied:

1. Prevent re-growth on potato dumps.
2. Destroy all groundkeepers.
3. Plant disease free seed.
4. Use generous soil ridges to protect tubers.

Integrate this approach with a fungicide programme.

## Early Crops

In first and second early potato crops, particularly those grown in the same locality as main crop potatoes, an adequate and full blight protection programme should be applied right up to harvesting or haulm desiccation. This will protect the early crop while helping to reduce disease risk to later crops.

## Maincrops

Disease prevention programmes require regular and season long fungicide use to limit foliar blight development. However, as an effective fungicide programme will preserve leaf area there may be more risk of infecting tubers at harvest, particularly during “heavy” blight years. Completion of the control programme should therefore include a complete haulm desiccant. Lifting of the crop should not take place for at least 10 days after **COMPLETE KILL** of the haulm. Crops intended for storage should not be lifted while there is any green tissue **AT ALL** on the leaves or stem bases.

## Blight Risk Assessment

The risk of disease is affected by **weather conditions** (during the crop life) and **crop location**:

**Weather Conditions** - Spread of disease occurs under warm, humid conditions. Preferably use a reliable decision support system to determine what frequency of fungicide treatment is appropriate and fungicide type required.

Note: Blight forecasting has often been based on the occurrence of “Smith periods”. A “Smith period” is a 48 hour period in which the minimum temperature is 10°C or more and the relative humidity exceeds 90% for at least 11 hours during the first 24 hours and for at least 11 hours again during the final 24 hours. However, any period of warm, humid weather increases blight risk.

**Crop Location** - Locations with the highest probability of blight problems are:

1. Areas of the country where extensive main crop or early production takes place e.g. East Anglia, the south west or the west.
2. Areas where climatic conditions that encourage disease development occur on a frequent basis e.g. The south west, the west and the Fens.

## BACKGROUND INTO ALTERNARIA CONTROL

The difenoconazole component of AMPHORE PLUS provides moderate activity against *Alternaria* spp. that cause Early blight. Primary infection can come into crops early in the rapid canopy growth phase. AMPHORE PLUS provides early protection against this and reduces the risk of secondary infection and disease progression later on when conditions become more favourable.

## TIMING

AMPHORE PLUS is a protectant fungicide and therefore the spray programme must start BEFORE blight enters the crop. Commence spraying at the first blight warning or when local weather conditions are favourable for the disease. *Alternaria* spray applications are best timed before visual symptoms are seen or at the very earliest signs of infection.

Intervals between applications of AMPHORE PLUS should be reduced as blight risk increases, so that **protection** of the crop can be maintained.

Applications of AMPHORE PLUS should be made at 7-10 day intervals depending on disease pressure. As disease pressure and the risk of late blight infection increase, the interval should be shortened.

## **Rates of Use**

Apply AMPHORE PLUS at 0.6 litres product per hectare. Up to three applications may be made per crop.

Applications of AMPHORE PLUS can be made up to 3 days before harvest.

## **MIXING AND SPRAYING**

### **Spray Volume**

Apply AMPHORE PLUS in 200-600 litres of water per hectare.

### **Spray Nozzles**

A medium quality spray\* is preferred for application of AMPHORE PLUS (\* as defined by The British Crop Protection Council). A spray pressure of 2 - 3 bar is recommended.

### **Mixing**

Make sure the sprayer is clean and set to give an even application at the correct volume.

Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of AMPHORE PLUS and continue agitation whilst adding the rest of the water.

Agitate the mixture thoroughly before use and continue agitation during spraying and any stoppages.

Thoroughly wash all spray equipment with water immediately after use.

Do not leave the spray liquid in the sprayer for long periods (such as during meal breaks or overnight).

## **PHYTOPHTERA RESISTANCE MANAGEMENT**

AMPHORE PLUS contains a CAA fungicide (FRAC code no. 40). To minimise the risk of resistance development in the pathogen population the following guidelines for CAA fungicides (based on an average number of 12 fungicide applications/season for blight control) should be followed:-

1. Where possible, use formulated/tank mixtures or an alternating strategy using fungicides from different mode of action groups.
2. Where CAA fungicides are applied as a mixture (co-formulated or as a tank mix) up to six applications (or max. of 50% of the total number of applications) may be made per crop or season.
3. Alternatively, where CAA fungicides are applied alone (without a co-formulant or tank mix partner) a maximum of four applications (or 33% of the total number of applications) may be made per crop or season.
4. No more than 3 applications of any CAA fungicide should be made consecutively.
5. Further information on suitable tank mix products and resistance management strategies is available from FRAG and BPC websites.

## **ALTERNARIA RESISTANCE MANAGEMENT**

AMPHORE PLUS contains a DMI fungicide (FRAC code no. 3). To minimise the risk of resistance development in the pathogen population to AMPHORE PLUS the following guidelines should be followed:-

1. Where possible use an alternating strategy using fungicides from different mode of action groups that must be active against the pathogen.

2. Further information on suitable tank mix products and resistance management strategies is available from FRAG and BPC websites.

**This product is to be used only in accordance with the recommendations and instructions given on the label provided with this pack.**

**To access the Safety Data Sheet for this product, scan QR code:**



**Alternatively, contact your supplier**