



# Evagio<sup>®</sup> Forte

syngenta.

**GROUP 21 | 40 FUNGICIDES**



Product registration number: MAPP 20854  
UFT: 9PQ7-FOSV-000J-XCTR

A suspension concentrate fungicide containing  
100 g/L of amisulbrom and 200 g/L of mandipropamid.

For the control of Foliar blight (*Phytophthora infestans*)  
in potatoes.

*The (COSHH) Control of Substances Hazardous to Health  
Regulations may apply to the use of this product at work.*

Syngenta UK Limited, Jealott's Hill International Research  
Centre, Bracknell, Berkshire, RG42 6EY  
Tel +44 (0) 1223 883400

**In case of toxic or transport emergency ring  
+44 (0) 1484 538444 (24hr)**

**STORE IN A COOL, DRY PLACE. PROTECT FROM FROST.**

## Safety Precautions

### (a) Operator protection

Engineering control of operator must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate. However engineering control may replace personal protective equipment if COSHH assessment shows that they provide an equal or higher standard of protection. KEEP OUT OF THE REACH OF CHILDREN.

Do not eat, drink or smoke when using the product.  
WASH HANDS AND EXPOSED SKIN before meals and after work.

# 5 litres



### (b) Environmental protection

To protect aquatic organisms, respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

**DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a horizontal boom sprayer either a LERAP must be carried out in accordance with CRD published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### (c) Storage and Disposal

KEEP IN ORIGINAL CONTAINER tightly closed, in a safe place. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely.

© Syngenta 2025


This product label is compliant with the CPA Voluntary Initiative (VI) guidance.



Product names marked © or ™, the ALLIANCE FRAME, the SYNGENTA Logo and the PURPOSE ICON are Trademarks of a Syngenta Group Company

L1131428 GBRI/01D PPE 4224283

A suspension concentrate containing 100 g/L of amisulbrom and 200 g/L of mandipropamid.



### Warning

**Causes serious eye irritation.**

**Suspected of causing cancer.**

**Very toxic to aquatic life with long lasting effects.**

Obtain special instructions before use.

Wear protective gloves/eye protection/face protection.

IF exposed or concerned: get medical advice/attention.

If eye irritation persists: get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

**To avoid risks to human health and the environment comply with the instructions for use.**

**MAPP 20854 UFI: 9PQ7-F0SV-000J-XCTR**

### IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crop	Maximum individual dose (L/ha)	Maximum total dose (L/ha)	Maximum number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Potato	0.6	1.8	3	6	7 days before harvest

**READ LABEL BEFORE USE: USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE: FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS**

This leaflet is part of the approved Product Label.

## **DIRECTIONS FOR USE**

**IMPORTANT:** This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

EVAGIO® FORTE can be used on all varieties of potatoes including seed potatoes.

## **PROPERTIES OF EVAGIO FORTE**

EVAGIO FORTE is a foliar fungicide for the preventative control of foliar blight (*Phytophthora infestans*) in potatoes and is highly active against spore germination. Uptake into the leaf tissues assures good translaminar and local systemic activity and inhibits mycelial growth during the incubation period. By reducing foliar blight, EVAGIO FORTE can also reduce the incidence of tuber blight.

## **DISEASE CONTROLLED**

Control of foliar blight (*Phytophthora infestans*) in potatoes.

## **BACKGROUND TO POTATO BLIGHT CONTROL**

Foliar blight (*Phytophthora infestans*) is potentially a devastating disease of potatoes. In commercial production, a season long disease prevention policy is essential. First of all ensure that other control methods are being satisfactorily applied:

1. Prevent re-growth on potato dumps.
2. Destroy all groundkeepers.
3. Plant disease free seed.
4. Use generous soil ridges to protect tubers.

Integrate this approach with a fungicide programme:

### **Early Crops**

In first and second early potato crops, particularly those grown in the same locality as main crop potatoes, an adequate and full blight protection programme should be applied right up to harvesting or haulm desiccation. This will protect the early crop while helping to reduce disease risk to later crops.

### **Maincrops**

Disease prevention programmes require regular and season long fungicide use to limit foliar blight development. However, as an effective fungicide programme will preserve leaf area there may be more risk of infecting tubers at harvest, particularly during

“heavy” blight years. Completion of the control programme should therefore include a complete haulm desiccant. Lifting of the crop should not take place for at least 10 days after COMPLETE KILL of the haulm. Crops intended for storage should not be lifted while there is any green tissue AT ALL on the leaves or stem bases.

### **Blight Risk Assessment**

The risk of disease is affected by weather conditions (during the crop life) and crop location:

#### **Weather Conditions**

Spread of disease occurs under warm, humid conditions. Preferably use a reliable decision support system to determine what frequency of fungicide treatment is appropriate and fungicide type required.

#### **Crop Location**

Locations with the highest probability of blight problems are:

1. Areas of the country where extensive main crop or early production takes place e.g. East Anglia, the south west or the west.
2. Areas where climatic conditions that encourage disease development occur on a frequent basis e.g. The south west, the west and the Fens.

### **TIMING**

EVAGIO FORTE is a protectant fungicide so following good agricultural practice the programme should start BEFORE blight enters the crop. Commence spraying at the first blight warning or when local weather conditions are favourable for the disease.

Intervals between applications of EVAGIO FORTE should be reduced as blight risk increases, so that protection of the crop can be maintained. Applications of EVAGIO FORTE should be made at 7-10 day intervals depending on disease pressure. As disease pressure and the risk of foliar blight infection increase, the interval should be shortened.

### **Rates of Use**

Apply EVAGIO FORTE at 0.6 L product per hectare. Up to three applications may be made per crop between BBCH 51 and 89. For non-flowering varieties, applications can commence from complete row closure, as first individual buds (1–2 mm) of first inflorescence visible (main stem) will normally coincide with growth stage. Allow a minimum of 6 days between applications. Applications of EVAGIO FORTE can be made up to 7 days before harvest.

## **MIXING AND SPRAYING**

### **Spray Volume**

Apply EVAGIO FORTE in a recommended 200-400 litres of water per hectare.

### **Spray Nozzles**

A medium quality spray\* is preferred for application of EVAGIO FORTE (\* as defined by The British Crop Protection Council). A spray pressure of 2 - 3 bar is recommended.

### **Mixing**

Make sure the sprayer is clean and set to give an even application at the correct volume. Fill the spray tank with half the required volume of clean water and start agitation. Add the required amount of EVAGIO FORTE and continue agitation whilst adding the rest of the water. Agitate the mixture thoroughly before use and continue agitation during spraying and any stoppages.

Thoroughly wash all spray equipment with water immediately after use. Drain the system completely and rinse spray tank, boom and nozzles three times with clean water until the foam and all traces of product have been removed. Do not leave the spray liquid in the sprayer for long periods (such as during meal breaks or overnight).

## **RESISTANCE MANAGEMENT**

EVAGIO FORTE contains mandipropamid, a CAA fungicide (FRAC code no. 40) and amisulbrom, a Qi-site inhibitor (Qil) fungicide (FRAC code no. 21). To minimise the risk of resistance development in the pathogen population the following guidelines for blight fungicides (based on an average number of 12 fungicide applications/season for blight control) should be followed:


1. Where possible, use an alternating strategy using fungicides from different mode of action groups.
2. Where CAA fungicides are applied as a mixture (co-formulated or as a tank mix) up to six applications (or max. of 50% of the total number of applications) may be made per crop or season.
3. No more than 2 applications of any CAA fungicide should be made consecutively.
4. Further information on suitable tank mix products and resistance management strategies is available from FRAG and BCPC websites.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier.

A suspension concentrate containing 100 g/L of amisulbrom and 200 g/L of mandipropamid.



### Warning

**Causes serious eye irritation.**

**Suspected of causing cancer.**

**Very toxic to aquatic life with long lasting effects.**

Obtain special instructions before use.

Wear protective gloves/eye protection/face protection.

IF exposed or concerned: get medical advice/attention.

If eye irritation persists: get medical advice/attention.

Collect spillage.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazol-3(2H)-one. May produce an allergic reaction.

**To avoid risks to human health and the environment comply with the instructions for use.**

**MAPP 20854 UFI: 9PQ7-F0SV-000J-XCTR**

### IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL FUNGICIDE

Crop	Maximum individual dose (L/ha)	Maximum total dose (L/ha)	Maximum number of treatments (per crop)	Minimum spray interval (days)	Latest time of application
Potato	0.6	1.8	3	6	7 days before harvest

**READ LABEL BEFORE USE: USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE: FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS**