



# Kayak<sup>®</sup>

syngenta<sup>®</sup>

**GROUP 9 FUNGICIDE**



**Product registration number:** MAPP 14847 **UFI:** XKX5-HOP0-C00U-86QA

An emulsifiable concentrate formulation containing 300 g/l cyprodinol.

For use against eyespot, net blotch, powdery mildew and *Rhynchosporium* on winter and spring barley.

## SAFETY PRECAUTIONS

### (a) Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: **WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES** when handling the concentrate.

However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

**WASH SPLASHES** from skin and eye immediately.

**WASH HANDS AND EXPOSED SKIN** before eating, drinking or smoking and after work.

### (b) Environmental protection

To protect aquatic organisms, respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

**DO NOT ALLOW DIRECT SPRAY** from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within

1 m of the top of a ditch which is dry at the time of application. **DO NOT ALLOW DIRECT SPRAY** from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing waterbody. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme.



Before each spraying operation from a horizontal boom sprayer or broadcast air assisted sprayer either a LERAP must be carried out in accordance with CRD published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### (c) Storage and disposal

**KEEP IN ORIGINAL CONTAINER**, tightly closed, in a safe place.

**DO NOT RE-USE CONTAINER FOR ANY PURPOSE.**

**RINSE CONTAINER THOROUGHLY** by using an integrated pressure rinsing device or manually rinsing 3 times. Add washings to sprayer at time of filling and dispose of safely.

Syngenta UK Limited

Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY  
Tel +44 (0) 1223 883400

**In case of toxic or transport emergency ring +44 (0) 1484 538444 any time**

*The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.*

**PROTECT FROM FROST.**

**STORE IN A COOL, DRY PLACE.**

© Syngenta AG, 2025

This product label is compliant with the  
CPA Voluntary Initiative (VI) guidance.



The  
Voluntary  
Initiative

L1131571 GBRI/01D PPE 4224440


# 5 litres

Product names marked © or ™, the ALLIANCE FRAME  
the SYNGENTA Logo and the PURPOSE ICON  
are Trademarks of a Syngenta Group Company



**KAYAK®**

Emulsifiable concentrate formulation containing 300 g/l cyprodinil.

**Warning**

**May cause an allergic skin reaction.**

**Very toxic to aquatic life with long lasting effects.**

Avoid breathing mist or vapours.

Wear protective gloves.

If skin irritation or rash occurs. Get medical advice/attention.

Take off contaminated clothing and wash it before reuse.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

**MAPP: 14847 UFI: XKX5-H0P0-C00U-86QA**

**IMPORTANT INFORMATION**

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE.

For use on Winter and Spring Barley.

Crops	Maximum individual dose (litres/product/ha)	Maximum number of treatments (per crop)	Latest time of application
Barley	1.5	2	Up to and including first awns visible stage (GS49)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

## **DIRECTIONS FOR USE**

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be carefully read in order to obtain safe and successful use of this product.

KAYAK® is an emulsifiable concentrate formulation containing 300 g/l cyprodinil for use as a fungicide on winter and spring barley.

## **DISEASES CONTROLLED**

Eyespot (*Pseudocercospora herpotrichoides*)

Powdery mildew (*Erysiphe graminis*)

Net blotch (*Pyrenophora teres*)

*Rhynchosporium secalis*

## **CROP SPECIFIC INFORMATION**

### **Crops**

All varieties of winter and spring barley.

### **Timing**

Treatment at early stages of disease development will ensure the best results. Further applications may be needed if disease attack is prolonged.

Eyespot: For best control of eyespot spray KAYAK before or during the period of stem extension in the spring (GS 30-32). Control of eyespot may be reduced when very dry conditions follow application.

Powdery Mildew, net blotch, *Rhynchosporium*: Application at first signs of disease is recommended.

For Powdery mildew control use early in the spray programme to make best use of the protectant activity. Use with an eradicant partner such as Corbel, Opus, Proline, Fandango or Helix for established infections.

For Net Blotch and *Rhynchosporium* control KAYAK should be used in tank mixture with another partner with a different mode of action such as AMISTAR, AMISTAR OPTI, CHEROKEE, Fandango, Helix, Opus and Proline.

### **Rates of Use**

Apply 1.5 litres of product per hectare. With a maximum treatment of 3 litres product per hectare (2 treatments per crop).

The latest time of application for KAYAK is before first spikelet of inflorescence just visible stage.

## **MIXING AND SPRAYING**

Make sure the sprayer is set to give an even application at the correct volume. In all cases add KAYAK to the spray tank first, followed by any additional products. Fill the spray tank with half the required volume of clean water and start agitation.

Add the required amount of KAYAK to the spray tank. Agitate the mixture thoroughly before use and continue agitation during spraying. Thoroughly wash all spray equipment with water immediately after use.

Wash out containers thoroughly, preferably using an integrated pressure rinsing device, or manually rinse three times.

Add washings to the sprayer at the time of filling. Complete filling to the required volume and continue to agitate throughout the spraying operation.

Do not leave the spray liquid in the sprayer for long periods (such as during meal breaks or overnight).

### **Spray Application**

Apply KAYAK in 200 litres of water per hectare using a tractor-mounted sprayer and a medium spray as defined by BCPC.

A spray pressure of 2-3 bars is recommended.

Thoroughly wash all spray equipment with water after use.

## **RESISTANCE MANAGEMENT**

KAYAK contains the aniline-pyrimidine fungicide cyprodinil. To reduce the risk of diseases developing resistance to cyprodinil, KAYAK should be used as part of a resistance management strategy that includes fungicides with a different mode of action.

For specific advice please refer to timing section in CROP SPECIFIC INFORMATION section above.

## **COMPATIBILITY**

KAYAK is compatible with AMISTAR, AMISTAR OPTI, CHEROKEE, Fandango, Helix, Opus and Proline.

## **SAFETY PRECAUTIONS**

### **(a) Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

**WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES** when handling the concentrate.

However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

**WASH SPLASHES** from skin and eye immediately.

**WASH HANDS AND EXPOSED SKIN** before eating, drinking or smoking and after work.

### **(b) Environmental protection**

To protect aquatic organisms, respect an unsprayed buffer zone distance to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing waterbody, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application.

DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing waterbody. Aim spray away from water.

This product qualifies for inclusion within the Local Environmental Risk Assessment for Pesticides (LERAP) Scheme. Before each spraying operation from a horizontal boom sprayer or broadcast air assisted sprayer either a LERAP must be carried out in accordance with CRD published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for inspection for three years.

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### **(c) Storage and disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

DO NOT RE-USE CONTAINER FOR ANY PURPOSE.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing 3 times.

Add washings to sprayer at time of filling and dispose of safely.

To access the Safety Data Sheet for this product, scan QR code:




Alternatively, contact your supplier.





**KAYAK®**

Emulsifiable concentrate formulation containing 300 g/l cyprodinil.

**Warning**

**May cause an allergic skin reaction.**

**Very toxic to aquatic life with long lasting effects.**

Avoid breathing mist or vapours.

Wear protective gloves.

If skin irritation or rash occurs. Get medical advice/attention.

Take off contaminated clothing and wash it before reuse.

Collect spillage.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

**MAPP: 14847 UFI: XKX5-H0P0-C00U-86QA**

**IMPORTANT INFORMATION**

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE.

For use on Winter and Spring Barley.

Crops	Maximum individual dose (litres/product/ha)	Maximum number of treatments (per crop)	Latest time of application
Barley	1.5	2	Up to and including first awns visible stage (GS49)

**READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.**

