

Levancio

GROUP 3 FUNGICIDE

Product registration number: MAPP 20580

An emulsifiable concentrate (EC) fungicide containing 125 g/L prothioconazole and 125 g/L tebuconazole for the control of certain diseases in wheat, barley, oats, winter rye and oilseed rape.

In case of toxic or transport emergency ring +44 (0) 1484 538444 (24hr)

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Authorisation Holder	Marketing Company
Clayton Plant Protection Ltd., Unit F10, Bracetown Business Park, Clonee, Co. Meath. Ireland Tel: +353 1 8210127	Syngenta UK Ltd Jealott's Hill International Research Centre, Bracknell, Berkshire, RG42 6EY Tel: Cambridge (01223) 883400

**STORE IN A COOL, DRY PLACE.
PROTECT FROM FROST.**



SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when applying the product.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT DRINK OR SMOKE.

WASH ANY CONTAMINATION from eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

IF YOU FEEL UNWELL, seek medical advice (show label where possible).

Environmental Protection

Do NOT CONTAMINATE SURFACE WATERS OR DITCHES with the product or its container. (Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads).



DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environmental Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.

Storage and Disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP IN ORIGINAL CONTAINER tightly closed in a safe place.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDINGSTUFFS.

PROTECT FROM FROST.



**The
Voluntary
Initiative**

This product label is compliant with the
CPA Voluntary Initiative (VI) guidance.

5 litres

Product names marked © or ™, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company

Levancio

Contains 125 g/L prothioconazole, 125 g/L tebuconazole.
Also contains N,N,-dimethyl decanamide.

Warning

Causes skin irritation.

May cause an allergic skin reaction.

May cause respiratory irritation.

Suspected of damaging the unborn child.

Toxic to aquatic life with long lasting effects.

Keep out of reach of children.

Wear protective gloves/protective clothing/eye protection/face protection.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.

Collect spillage

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for triple rinsed empty clean containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



MAPP 20580

IMPORTANT INFORMATION: ONLY FOR USE AS A PROFESSIONAL FUNGICIDE

Crop	Maximum individual dose (L/ha)	Maximum number of treatments per crop	Latest time of application
Rye (winter), wheat	1.0	2	End of flowering (BBCH 69)
Barley, oats	1.0	2	Up to beginning of flowering (BBCH 61)
Oilseed rape	1.0	2	56 days before harvest

Other specific restrictions: A minimum of 21 days must be observed between applications.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Levancio contains prothioconazole and tebuconazole, both triazole fungicides, recommended for control of a wide range of diseases on certain cereals and winter oilseed rape.

CROPS

Levancio may be used on all commercial varieties of winter and spring barley, winter and spring wheat, winter rye, winter and spring oats and winter oilseed rape.

RATE OF USE

Apply Levancio at 1.0 L/ha.

DISEASE CONTROLLED

Cereals

	Wheat	Barley	Oats	Rye
Septoria leaf blotch	C			
Glume blotch	C			
Yellow rust	C	C		
Brown rust	C	C		C
Powdery Mildew	MC	C	C	C
Crown rust			C	
Rhynchosporium		C		C
Net blotch		C		
Eyespot	reduction	reduction	C	reduction
Tan Spot	C			
Fusarium Ear Disease complex	reduction	reduction		
Sooty moulds	reduction	reduction		

C = control MC – moderate control

APPLICATION TIMING

A minimum interval of 21 days must be observed between applications in all crops.

CEREALS

Eyespot (*Tapesia spp.*)

Spray in the spring at the first sign of disease, from when the leaf sheaths begin to become erect until the 2nd node is detectable (GS 30-32).

Septoria Leaf Blotch and Glume Blotch (*Septoria tritici* and *Stagonospora nodorum*)

Apply before disease is established in the crop. To protect the upper leaves and ear apply Levancio at full flag leaf emergence (GS 37) up to mid-flowering (GS 65). Where disease pressure remains high application may be repeated. Applications to upper leaves where *S. tritici* symptoms are present are likely to be less effective. Levancio contains two DMI fungicides. Resistance to some DMI fungicides has been identified in Septoria leaf blotch (*Mycosphaerella graminicola*) which may seriously affect the performance of some products. For further advice on resistance management in DMIs contact your agronomist or specialist advisor, and visit the FRAG-UK website.

Powdery Mildew (*Blumeria graminis*)

Apply Levancio at the first signs of disease. Where disease pressure remains high application may be repeated. Levancio will only provide moderate control of powdery mildew in wheat.

Yellow Rust (*Puccinia striiformis*)

Apply Levancio at the first signs of disease. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

Brown Rust (*Puccinia recondita* and *P horde*)

Apply Levancio at the first signs of disease. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

Crown Rust (*Puccinia coronata*)

Apply Levancio at the first signs of disease. Levancio controls crown rust in winter and spring oats. A second application may be made later if re-infection occurs. Applications made to established infections are likely to be less effective.

Tan Spot (*Pyrenophora tritici-repentis*)

Apply Levancio at the first signs of disease in spring or early summer. Where disease pressure remains high application may be repeated.

Ear Disease Complex

Apply Levancio soon after ear emergence until the end of flowering (GS 59-69) for control of Fusarium ear blight (or moderate control in barley) and reduction of sooty moulds. Control of ear diseases can result in cleaner, brighter ears. Through the control of ear blight, Levancio effectively reduces the level of the Fusarium mycotoxin deoxynivalenol (DON) in wheat grain. However, where Fusarium levels are high, the reduction achieved may not always be sufficient to ensure that DON levels fall below the statutory limit.

Leaf Blotch (*Rhynchosporium secalis*)

Apply Levancio in spring at the first signs of disease. For severe infections a second application may be necessary later.

Net Blotch (*Pyrenophora teres*)

Apply Levancio in autumn at the first signs of disease in spring/early summer. For severe infections, a second application later will give most effective control when conditions remain favourable for disease development.

OILSEED RAPE

Light Leaf Spot (moderate control)

Apply Levancio protectively in autumn/winter (usually late October to early December). Follow up treatment(s) may be required in early spring from the onset of stem elongation, depending on disease development.

Phoma Leaf spot/Stem Canker

Apply Levancio in autumn at the first sign of disease. Repeat treatment in late autumn/winter, if disease symptoms reoccur.

Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Apply Levancio at early to full flower.

RESISTANCE

Repeated application of Levancio alone should not be used on the same crop. Tank-mixtures or alternation with fungicides having a different mode of action have been shown to protect against the development of resistant forms of disease. The possible development of disease strains resistant to Levancio cannot be excluded or predicted. Where such resistant strains occur, Levancio is unlikely to give satisfactory control. Strains of Light Leaf Spot resistant to azole fungicides are known to exist. To avoid development of resistance apply product protectively

in response to disease forecasts. Where possible, when Light Leaf Spot is present, avoid the use of azole based fungicides when targeting other diseases such as Sclerotinia at mid flowering.

APPLICATION METHOD

Spray equipment should be thoroughly cleaned before use. Apply with tractor mounted/trailed sprayer in 100-400 litres of water per hectare. The higher volume is recommended on dense crops to ensure good coverage and penetration. Apply as a MEDIUM quality spray (as defined by BCPC). A spray pressure of 2-3 Bar is recommended.

MIXING

Thoroughly shake the pack before use. Add the required quantity to the half-filled tank with the agitation system in operation and then fill to the required level. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray equipment should be cleaned immediately after use. Sprayers should be thoroughly cleaned with water and detergent after use, and filters and jets checked for damage and blockages.

To access the Safety Data Sheet for this product, scan QR code:



Alternatively, contact your supplier

Levancio

Contains 125 g/L prothioconazole, 125 g/L tebuconazole.
Also contains N,N,-dimethyl decanamide.

Warning

Causes skin irritation.

May cause an allergic skin reaction.

May cause respiratory irritation.

Suspected of damaging the unborn child.

Toxic to aquatic life with long lasting effects.

Keep out of reach of children.

Wear protective gloves/protective clothing/eye protection/face protection.

IF exposed or concerned: Call a POISON CENTRE or doctor/physician.

Collect spillage

Dispose of contents/container to a licensed hazardous-waste disposal contractor
or collection site except for triple rinsed empty clean containers which can be disposed
of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.



MAPP 20580

IMPORTANT INFORMATION: ONLY FOR USE AS A PROFESSIONAL FUNGICIDE

Crop	Maximum individual dose (L/ha)	Maximum number of treatments per crop	Latest time of application
Rye (winter), wheat	1.0	2	End of flowering (BBCH 69)
Barley, oats	1.0	2	Up to beginning of flowering (BBCH 61)
Oilseed rape	1.0	2	56 days before harvest

Other specific restrictions: A minimum of 21 days must be observed between applications.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.