

United Kingdom

Published on Syngenta (<https://www.syngenta.co.uk>)

[Home](#) > WAKIL XL

WAKIL XL



WAKIL XL

Last updated:

16.08.2018

Seed Treatment

Crop protection

Authorisation Number:

MAPP/MAFF No.: 17217

Pack size:

300 kg

Composition:

100 g/kg

cymoxanil

50 g/kg

fludioxonil

175 g/kg

metalaxyl-M

Chemical Family:

Cyanoacetamide-oxime

Phenylpyrrole

Phenylamide

Activity Group:

FRAC Code 27 (cymoxanil), Group E2 FRAC code 12 (fludioxonil), Group A1 FRAC code 4 (metalaxyl-M)

NANUM number:

NAN2371/2015

Formulation:

WG (Water Dispersible Granule)

WAKIL XL is a co-formulation of cymoxanil, fludioxonil and metalaxyl-M for the control of downy mildew, *Pythium* spp. and *Ascochyta* spp. in combining and vining peas.

Product CP: Tabs

- [Registrations](#)
- [Application Advice](#)
- [Emergency](#)

[Registrations](#)

[WAKIL XL Peas](#)

Peas

Resistance management / status:

A minimum of at least 5 years between pea (or bean) crops is advised in fields known to be infested with downy mildew. Avoid drilling peas very early in the season when soil temperatures are lower. Avoid deep drilling.

Ascochyta

Maximum individual dose	Maximum total dose	Maximum number of applications	Pre Harvest Interval (PHI)	Aquatic Buffer Zone	Non Target Arthropods Buffer Zone	Non Target Plant Buffer Zone
2.0 kg/tonne		1	Before drilling	Unclassified	None	None
Stage of use:	Before drilling					
Restriction of Use:	Treated seed must not be used for food or feed. Sacks containing treated seed must not be re-used for food or feed. Treated seed must not be applied from the air. Treated seed must be labelled using the printed sacks, labels or bag tags supplied. Treated seeds must not be sown between 29 September and 1 April.					

Downy Mildew (moderate)

Maximum individual dose	Maximum total dose	Maximum number of applications	Pre Harvest Interval (PHI)	Aquatic Buffer Zone	Non Target Arthropods Buffer Zone	Non Target Plant Buffer Zone
2.0 kg/tonne		1	Before drilling	Unclassified	None	None

Stage of use: Before drilling
Restriction of Use: Treated seed must not be used for food or feed. Sacks containing treated seed must not be re-used for food or feed. Treated seed must not be applied from the air. Treated seed must be labelled using the printed sacks, labels or bag tags supplied. Treated seeds must not be sown between 29 September and 1 April.

Pythium spp.

Maximum individual dose	Maximum total dose	Maximum number of applications	Pre Harvest Interval (PHI)	Aquatic Buffer Zone	Non Target Arthropods Buffer Zone	Non Target Plant Buffer Zone
2.0 kg/tonne		1	Before drilling	Unclassified	None	None

Stage of use: Before drilling
Restriction of Use: Treated seed must not be used for food or feed. Sacks containing treated seed must not be re-used for food or feed. Treated seed must not be applied from the air. Treated seed must be labelled using the printed sacks, labels or bag tags supplied. Treated seeds must not be sown between 29 September and 1 April.

Application Advice

Mixing and spraying

See label for application advice

Rainfastness

Not applicable

Emergency

- In the event of a Toxic or Transport Emergency in the UK please telephone 01484538444 at any time.