

United Kingdom

Published on Syngenta (<https://www.syngenta.co.uk>)

[Home](#) > MICARAZ

MICARAZ

MICARAZ

Last updated:

08.08.2018

Fungicide

Authorisation Number:

MAPP/MAFF No.: 16106

Composition:

90 g/l

epoxiconazole

125 g/l

isopyrazam

Chemical Family:

Triazole

SDHI

Activity Group:

Group G1 FRAC code 3

Group C2 FRAC code 7

Formulation:

SC (Suspension Concentrate)

MICARAZ is a co-formulation of epoxiconazole and isopyrazam for the control of a range of diseases in winter wheat, rye and triticale and both winter and spring barley

Product CP: Tabs

- [Registrations](#)
- [Application Advice](#)
- [Emergency](#)

Registrations

MICARAZ Barley

Barley

Resistance management / status:

Do not apply more than two foliar applications of SDHI-containing products to any crop.

MICARAZ Triticale

Triticale

Resistance management / status:

Do not apply more than two foliar applications of SDHI-containing products to any crop.

MICARAZ Winter Rye

Winter Rye

Resistance management / status:

Do not apply more than two foliar applications of SDHI-containing products to any crop.

MICARAZ Winter Wheat

Winter Wheat

Resistance management / status:

Do not apply more than two foliar applications of SDHI-containing products to any crop.

Microdochium leaf spot (moderate)

Maximum individual dose	Maximum number of applications	Pre Harvest Interval (PHI)	Aquatic Buffer Zone	Non Target Arthropods Buffer Zone	Non Target Plant Buffer Zone
1.0 l/ha	2	Before grain watery ripe stage (GS71)	LERAP B	None	None

Stage of use: From GS30 up to before grain watery ripe stage (GS71)

Restriction of Use: A minimum of 21 days must be observed between applications.

Tan spot (reduction)

Maximum individual dose	Maximum number of applications	Pre Harvest Interval (PHI)	Aquatic Buffer Zone	Non Target Arthropods Buffer Zone	Non Target Plant Buffer Zone
1.0 l/ha	2	Before grain watery ripe stage (GS71)	LERAP B	None	None

Stage of use: From GS30 up to before grain watery ripe stage (GS71)

Restriction of Use: A minimum of 21 days must be observed between applications.

Application Advice

Mixing and spraying

Half fill the sprayer with water, add the MICARAZ, allow to disperse before adding any other product then top up the tank whilst maintaining agitation. Apply using a medium quality spray using a spray pressure of 2-3 bar.

Water volume

100-400 l/ha

Rainfastness

1 hour

Emergency

- In the event of a Toxic or Transport Emergency in the UK please telephone 01484538444 at any time.