

This seed has been treated with

MAPP 14654



syngenta.

Contains metalaxyl-M and protects against Pythium spp. in a range of vegetable seed crops.

#### **SAFETY PRECAUTIONS**

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when handling treated seed.

To protect birds and wild mammals treated seed should not be left on the soil surface. Bury or remove spillages.

THIS SEED HAS BEEN CHEMICALLY TREATED.

Treated seed should be precision drilled.

If seed is present on the soil surface, or if spills have occurred, then, if conditions are appropriate, the field should be harrowed and then rolled to ensure good incorporation.

DO NOT HANDLE seed unnecessarily.

DO NOT USE TREATED SEED as food or feed.

DO NOT RE-USE SACKS THAT HAVE CONTAINED TREATED SEED for food or feed.

KEEP TREATED SEED SECURE from people, domestic stock / pets and wildlife at all times during storage and use.

WASH HANDS AND EXPOSED SKIN before meals and after work.

DO NOT APPLY TREATED SEED FROM THE AIR.

#### **NOTES**

1. Safe Handling of treated Seed

Avoid skin contact with treated seed and dust during all drilling operations. Launder coveralls daily.

2. Drilling

Check drill calibration before drilling for each batch of seed to ensure an accurate drilling rate. Avoid adverse seedbed conditions and deep or shallow drilling which may adversely affect crop establishment and reduce the level of pest control.

3. Storage

Seed should be stored in a cool, dry, well ventilated building and be drilled as soon as possible after treatment. Drill within the season of treatment.

4. Seed Spillages

In case of seed spillage, clean up as much as possible into the related seed sack and re-use the clean seed. Bury the remainder completely.

#### **Syngenta UK Limited**

CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE

Tel: Cambridge (01223) 883400

**In case of toxic or transport emergency ring +44 (0) 1484 538444 any time**

APRON<sup>®</sup>XL is a Registered Trademark of a Syngenta Group Company.

© Syngenta 2018

