



syngenta.

Contains Fludioxonil.

For the control of *Leptosphaeria maculans* in cabbage, *Alternaria spp.* in carrot, *Botrytis allii* in bulb onion and moderate control of *Colletotrichum dematium* in spinach.

SAFETY PRECAUTIONS

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the concentrate, or contaminated surfaces or treated seed.
WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) when bagging treated seed.

However engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

WHEN USING, DO NOT EAT, DRINK OR SMOKE.

TO PROTECT BIRDS TREATED SEED SHOULD NOT BE LEFT ON THE SOIL SURFACE. BURY OR REMOVE SPILLAGES.

Do not contaminate water with the product and its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

STORAGE

EMPTY CONTAINER COMPLETELY and dispose of safely. KEEP IN ORIGINAL CONTAINER tightly closed in a safe place. RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of safely. DO NOT RE-USE CONTAINER for any purpose.

Syngenta UK Limited

CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE
Tel: Cambridge (01223) 883400

**In case of toxic or transport emergency ring
+44 (0) 1484 538444 any time**

MAXIM® 480 FS is a Registered Trademark of a Syngenta Group Company.

© Syngenta 2018

