# **LUTIMOR TANK MIX LIST**

### GUIDANCE FOR TANK MIXING AND SEQUENCING WITH HORMONE HERBICIDES

- DO NOT mix LUTIMOR with products containing 2-4,D, MCPA or mecoprop.
- These hormone herbicides may be used in sequence with LUTIMOR, observing the following intervals:
  - o When LUTIMOR is applied first, leave a minimum of 7 days before applying hormone herbicides
  - o If hormone-containing products are applied first, leave 21 days before LUTIMOR is applied
- Halauxifen (Arylex), fluroxypyr or clopyralid products may be tank mixed with LUTIMOR when LUTIMOR is used at the **full label dose rate** for the target weed (refer to label).
- DO NOT MIX LUTIMOR with Halauxifen (Arylex), fluroxypyr or clopyralid products when targeting black-grass in barley.
- When targeting wild oats or ryegrass using supported doses of LUTIMOR, Halauxifen (Arylex), fluroxypyr or clopyralid products may be sequenced with LUTIMOR observing the same intervals indicated above.

# GUIDANCE FOR TANK MIXING AND SEQUENCING WITH SULFONYL-UREAS AND OTHER ALS INHIBITOR HERBICIDES

- Tank mixes with some broad-leaved weed specific sulfonyl-urea/ALS inhibitor herbicides can reduce grass weed control, particularly if LUTIMOR is applied at reduced rates.
- Only sulfonyl-urea/ALS inhibitor herbicides listed below may be tank mixed with LUTIMOR.
- DO NOT MIX LUTIMOR with SU's/ALS inhibitors when targeting black-grass in barley.
- Sulfonyl-urea/ALS inhibitor herbicides should only be tank mixed with LUTIMOR when LUTIMOR is used at the **full label dose rate** for the target weed (refer to label).
- When targeting <u>wild oats or ryegrass</u> using supported rates of LUTIMOR, in sequence with SUs/ALS inhibitors, observe the intervals indicated above for hormone herbicides.

The following tank mixes have been tested for physical compatibility with LUTIMOR at recommended rates of use and will mix in the sprayer tank. Syngenta UK Ltd will support 2 or 3-way tank mixes of LUTIMOR with any of the fungicides, herbicides, insecticides, PGR's or adjuvants listed. All other products listed have not been tested for crop safety or product performance, therefore use is at the user's own risk. For further information on compatibilities, contact Syngenta UK Ltd on 0800 169 6058.

### **FUNGICIDES**

**AMISTAR** Helix Prosaro Inception Xpro Protefin Ascra Xpro Aviator Xpro Jaunt1 **OUADRIS** Justice1 Caramba Revystar<sup>4</sup> Comet 200 KAYAK Rylox<sup>4</sup> KAVATUR PLUS<sup>1</sup> Cebara Siltra Xpro CERATAVO PLUS1  $Talius^1 \\$ LAMAST **ELATUS ERA** Librax UNIX

 $\begin{array}{ccc} ELATUS \ PLUS^1 & MIRROR & VELOGY \ PLUS^1 \\ Fandango^1 & Mobius & WEAVER + Scyon \end{array}$ 

Folicur PLEXEO Zephyr

Haven Proline 275

## **HERBICIDES**

 $\begin{array}{cccc} Ally \ Max \ SX^2 & Hiatus^3 & Renitar \\ Ergon^3 & Mattera & Spitfire \\ Finish \ SX & Pixxaro & Whorl \\ Harmony \ M \ SX & Presite \ SX & Zypar \end{array}$ 

#### **INSECTICIDES**

HALLMARK ZEON Decis Protech KUNG FU

# PGR's

3C Chlormequat 750 Cerone MODDUS

Adjust Cleancrop Alatrin Evo MODDUS + chlormequat

Canopy Medax Max Terpal

#### **ADJUVANTS**

If LUTIMOR is applied at the reduced rate of below 0.6 l/ha, an adjuvant should be added for optimum results. Syngenta will support the use of the adjuvants listed.

ADIGOR Phase II Toil
Nelson Sorrento<sup>5</sup> Zarado

Syngenta have not tested any other adjuvants, however, the use of other MSO's not included in the list above will be supported.

## TRACE ELEMENTS

Syngenta UK Ltd will support the tank mixing of **one** of the named trace elements with an existing 2 or 3-way tank mix including LUTIMOR. Syngenta UK Ltd advise that the trace element part of the tank mix is added to the sprayer tank last with constant agitation and the mixture is sprayed without delay.

BittersalzHeadland Multiple ProYaraVita Coptrel 500Headland Boron 15Headland TriceYaraVita Croplift ProHeadland CopperHeadland Zinc 7001YaraVita MancozinHeadland Manganese 15%YaraVita Bortrac 150YaraVita Mancuflo

### **NOTES**

Before using any tank mixture, consult and comply with the recommendations of the partner products. Each product should be added separately to the bulk of the water in the spray tank and thoroughly mixed before adding the next chemical. Always use constant agitation of the sprayer tank during mixing, transportation and application. Spray immediately.

- <sup>1</sup> Continuous agitation required.
- <sup>2</sup> Ally Max SX = can read across to all FMC SUs.
- <sup>3</sup> Tested and supported by Rotam.
- <sup>4</sup> Tested and supported by BASF.
- <sup>5</sup> Tested and supported by Interagro.

The following 3-way tank mixes have been tested and are **physically compatible**:

LUTIMOR + Inception Xpro + Mobius

LUTIMOR + Inception Xpro + Jaunt<sup>1</sup>

LUTIMOR + Inception Xpro + Zephyr

LUTIMOR + Rylox + Arizona<sup>4</sup>

LUTIMOR + Rylox + Sorrento<sup>4</sup>

LUTIMOR + Rylox + Arizona + Sorrento<sup>4</sup>

LERAP: None Rainfastness: 1 hour

# Use plant protection products safely. Always read the label and product information before use.

©Syngenta AG, February 2022



# For further information contact us on 0800 169 6058 e-mail: customer.services@syngenta.com

Syngenta UK Ltd, Fulbourn, Cambridge CB21 5XE.

Tel: (01223) 884300. Fax: (01223) 281245. Website: http://www.syngenta.co.uk