

# Environmental Information Sheet

## AMISTAR TOP MAFF/MAPP 12761



A SC formulation containing 200 g/litre azoxystrobin, a strobilurin fungicide, and 125 g/l difenoconazole, an azole fungicide, for use in carrots and leeks, Brussels sprouts, cabbage, cauliflower, broccoli/calabrese, kale (winter greens) and collards (spring greens).

Maximum application rate: 1 litres per hectare

Maximum no. of applications: 2 per crop

Section	Profile
<p><b>1. WILDLIFE</b></p> <p><b>Mammals and Birds</b></p>	<p>Amistar Top is not classified as <i>'Harmful to game, wild birds and animals'</i>.</p> <p>No risk management is necessary to protect wild mammals and birds. Amistar Top is of low toxicity to both mammals and birds. The risk to wild mammals and birds feeding on treated areas is low due to the limited amount of residues likely to be consumed.</p>
<p><b>2. BEES</b></p>	<p>No risk management is necessary. Amistar Top is of low risk to honey bees.</p>
<p><b>3. NON TARGET INSECTS AND OTHER ARTHROPODS</b></p>	<p>No risk management is necessary. Amistar Top poses a low risk to a range of arthropod species commonly found in and around treated fields.</p>
<p><b>4. AQUATIC LIFE</b></p>	<p>Amistar Top is classified as <i>'Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment.'</i></p> <p>No risk management is necessary. Amistar Top is highly toxic to fish and aquatic life. However, in practice the risk to aquatic life is low if used as recommended. Care should be taken to ensure that surface waters or ditches are not contaminated with the product or used container.</p> <p>Amistar Top is not categorised under the LERAP scheme.</p>

# Environmental Information Sheet

## AMISTAR TOP MAFF/MAPP 12761



Section	Profile
<p><b>5. SOIL and GROUNDWATER</b></p> <p><b>Earthworms</b></p> <p><b>Soil Micro-organisms</b></p>	<p>Difenoconazole is very persistent in soil. It is strongly absorbed to soil and not considered as mobile. Azoxystrobin is moderately persistent, and has moderate mobility in soil. Use of Amistar Top according to the label instruction presents a low risk to groundwater.</p> <p>Amistar Top is of moderate toxicity to earthworms. No risk management is necessary as the risk to earthworms is low.</p> <p>Amistar Top is unlikely to have any long-term effect on soil microbial activity. The risk is therefore considered to be low. No risk management is necessary.</p>
<p><b>6. NON-TARGET PLANTS</b></p>	<p>No effects on non-target plant species are expected when Amistar Top is used as recommended.</p>

**USE PLANT PROTECTION PRODUCTS SAFELY. ALWAYS READ THE LABEL AND PRODUCT INFORMATION BEFORE USE.**

Care must be taken to minimise the risk of surface water contamination from farmyard and field sources.

For further information about the environmental profile of this product contact:-

Company: Syngenta UK Ltd  
 Address: CPC4, Capital Park, Fulbourn, Cambridge CB21 5XE  
 Telephone: 0800 1696058  
 Fax: 01223 882195  
 e-mail: [customer.services@syngenta.com](mailto:customer.services@syngenta.com)

This Environmental Information Sheet was prepared in accordance with CPA Guidance Notes Version 4.

© copyright of Syngenta  
 The Amistar Top Trademark is the property of a Syngenta Group Company